

PHASE 1
BIRTH TO 20 YEARS



TOTAL CORRECTION EXPECTED.
6 MONTHS TO 1 1/2 YEARS OF CARE.
LIFETIME MAINTENANCE.
CARE RECOMMENDED.

DECREASED CURVE.
CHRONIC MISALIGNMENTS.
NORMAL BONE AND DISCS.
SOFT TISSUE CHANGES.

LOWERED RESISTANCE.
VAGUE AND INTERMITTENT
SYMPTOMS BECOMING ACUTE.

JOINT, DISC AND BONY TISSUE
CHANGES SIGNIFICANT
FIBROSIS.

ACUTE FUNCTIONAL
DISTURBANCES BECOMING
CHRONIC.

PHASE 2
20 TO 40 YEARS



SIGNIFICANT IMPROVEMENT OVER
A LONG TERM BASIS.
1 1/2 TO 2 1/2 YEARS OF CARE.
LIFETIME MAINTENANCE.

PHASE 3
40 TO 65 YEARS



USUALLY SLOW DOWN AND
OCCASIONALLY STOP THE
SUBLUXATION DEGENERATION
2 1/2 TO 3 1/2 YEARS OF CARE.
LIFETIME MAINTENANCE.

MAJOR DEGENERATIVE
CHANGES. LOSS OF BONE
MASS. VERTEBRAL COLLAPSE.
BEGINNING OF FUSION.

CHRONIC AND SEVERE HEALTH
PROBLEMS.

ALL CHANGES SEEN IN
PREVIOUS PHASES INCLUDING
SERIOUS BONE FUSIONS,
SPINAL COLLAPSE, AND
PERMANENT SUBLUXATIONS.

PERMANENT LACK OF
FUNCTION AND ILL HEALTH.
ADVANCING COMPLICATIONS.

PHASE 4
65 YEARS & OLDER



CARE GIVEN FOR RELIEF OF
SYMPTOMS. (NOT CORRECTION)
MUST WORK WITH AREAS OF SPINE
NOT AS SEVERELY AFFECTED.
LIFETIME CARE REQUIRED.